

BOSTON PUBLIC LIBRARY



3 9999 06317 688 5

49.15/3:

141

# BURNING DOVE STATUS REPORT

## 1970

Boston Public Library,  
Superintendent of Documents

SEP 15 1971

DEPOSITORY



UNITED STATES DEPARTMENT OF THE INTERIOR  
FISH AND WILDLIFE SERVICE  
BUREAU OF SPORT FISHERIES AND WILDLIFE  
Special Scientific Report — Wildlife No. 141



UNITED STATES DEPARTMENT OF THE INTERIOR  
Fish and Wildlife Service  
Bureau of Sport Fisheries and Wildlife

# MOURNING DOVE STATUS REPORT, 1970

*Compiled by*

*James L. Ruos  
Migratory Bird Populations Station  
Division of Wildlife Research*



Bureau of Sport Fisheries and Wildlife  
Special Scientific Report--Wildlife No. 141  
Washington, D.C. May 1971



# CONTENTS

Abstract .....	iv
Introduction .....	1
Procedure	
The Call-Count Survey .....	2
Quality checks of field data .....	2
Randomization of call-count routes .....	3
Physiographic stratification of call-count routes .....	3
Breeding Density Index .....	6
Determination of short-term population changes, 1969 to 1970 ...	8
Determination of long-term population trends by State, 1960 to 1970 .....	8
Determination of long-term trends by physiographic region, 1965 to 1970 .....	8
Computer analysis of dove call-count data .....	9
Statistical evaluation of data .....	9
Determination of population distribution .....	9
Findings	
Status of the United States dove population	
1970 population distribution .....	10
1969 to 1970 population changes .....	10
1960 to 1970 long-term population trends .....	10
Status of the Eastern Management Unit population	
1970 population distribution .....	15
1969 to 1970 population changes .....	15
1960 to 1970 long-term population trends .....	15
Status of the Central Management Unit population	
1970 population distribution .....	22
1969 to 1970 population changes .....	22
1960 to 1970 long-term population trends .....	22
Status of the Western Management Unit population	
1970 population distribution .....	23
1969 to 1970 population changes .....	23
1960 to 1970 population trends .....	23
Statistical significance of data	
1969 to 1970 population changes .....	23
1960 to 1970 long-term population trends .....	24
Acknowledgements .....	24
References .....	25
Tables .....	27

## ABSTRACT

Mourning dove population indexes increased by 4 percent from 1969 to 1970 in the Eastern Management Unit, but declined 2 percent in the Central Management Unit and 11 percent in the Western Management Unit. The changes were not statistically significant. The 1970 indexes were below the 10-year means, 1960-69, by 3 percent in the Eastern Unit, 15 percent in the Central, and 27 percent in the Western. Regression analyses of the call-count data for 1960-70 indicate a statistically significant downward trend in dove breeding populations in all management units; mean rates of decline were 1 percent a year in the Eastern Unit, 3 percent a year in the Central, and 4 percent a year in the Western.

Changes in the population indexes are described by State and physiographic region. The 1970 indexes were generally lower than in 1969 in the northern and western areas of the country, and generally higher in sections of the Rocky Mountains, the middle States, and the eastern seaboard. Regression analyses of 11 years' data, 1960 to 1970, indicate a statistically significant downward trend in population in much of the area represented by the northern Great Plains and Pacific border and southern border States. Trends are significantly upward in several scattered mid-latitude States.

## INTRODUCTION

Management of mourning doves in the United States is essentially the regulation of hunting to achieve proper harvest. The Call-Count Survey, conducted annually since 1953 by Federal, State, and independent observers on more than 800 prescribed routes, provides population data on which wildlife administrators rely in setting annual regulations. This report describes the methods employed to obtain and analyze those data and presents the status of the 1970 mourning dove breeding population.

Two versions of the dove status report, one preliminary and one final, are prepared annually. In 1970 the preliminary version was mailed to members of the Dove Regulations Committee a week before the regulations meeting in June at Washington, D.C. This timely distribution was made possible by the promptness of cooperators who sent their data directly to the Migratory Bird Populations Station immediately after completion of their surveys. The present report is the final version and contains additional survey data received too late for use in the preliminary version. As is customary, it will be distributed to all cooperators and will be available to interested organizations and individuals.

Basic data gathering and analyzing procedures used in this report were the same as those used in 1969 (Ruos, 1970).



## PROCEDURE

### The Call-Count Survey

Field studies have demonstrated the feasibility of the Call-Count Survey as a method for detecting annual changes in mourning dove breeding populations (Foote and Peters, 1952). Since 1953, these surveys have been conducted throughout the United States over a system of more than 800 established routes. Each call-count route has twenty, 3-minute listening stations spaced at 1-mile intervals, usually on lightly traveled secondary roads.

Each route is checked once between May 20 and June 10. Intensive studies in the eastern United States (Foote and Peters, 1952) indicated that dove calling is relatively stable during this period. Call-count surveys are not made when wind velocities exceed 12 miles per hour or when it is raining.

Records are kept of all doves seen or heard calling along the routes. The numbers heard calling during the 3-minute listening periods are totaled for each route to provide the data for determining the population index. The numbers of calls per dove and of doves seen are not currently used in the index calculation, but they are recorded. A detailed analysis of these and other pertinent data from past call counts is currently under study by the Migratory Bird Populations Station.

Studies by Frankel and Baskett (1961) and Jackson and Baskett (1964) have shown that unmated males call at a greater rate than mated males. This suggests that the reliability of the annual call-count census is reduced by the variability in the ratio of mated to unmated males. However, Wight (1964) observed that variations in the ratio of mated to unmated males, where the adult sex ratio approached equality, did not significantly alter the reliability of the dove call count for measuring annual trends of breeding mourning doves. Irby (1964) also found no evidence on his study area in Arizona that the numbers of unmated males materially affected call-count results.

### Quality checks of field data

Survey reports were examined to determine circumstances affecting the accuracy with which the routes were run and the data recorded. Records for routes run under unacceptable conditions were not analyzed. Reports on routes completed under the prescribed conditions but containing discrepancies or errors, or lacking data, were examined to ascertain whether parts were acceptable. If so, they were used in



analyses for which they were applicable. Where there was a change in observers on a route from one year to the next, the data were examined to determine whether an unexpected population change was apparent. When such differences exceeded those of the prescribed limits, they were attributed to differences in observers, and the data were not used in the current analyses.

#### Randomization of call-count routes

The original call-count routes (established between 1951 and 1956, and hereafter designated "management routes") were, in many instances, selected in areas of high-density dove populations and were not representative of populations over entire States or management units.

Randomly located routes were first employed in seven southeastern States in 1957 (Foote, Peters, and Finkner, 1958). A study of the random and management route data from these States confirmed earlier assumptions that a revision of the nationwide call-count survey routes should be undertaken if representative dove population indexes were to be obtained. This recommendation prompted the gradual selection and establishment of 912 randomly located call-count routes. In 1970, for the first time, random routes were established in all 48 conterminous States.

Both types of routes were run during the year of transition from management to random routes. This procedure permitted a direct comparison of data (Foote, Peters, and Finkner, 1958).

#### Physiographic stratification of call-count routes

Biologists recognize the limitation of sampling wildlife populations by political units. Census data collected and analyzed by ecological divisions represent better statistical design and could be expected to provide more precise information with the same effort.

An ecological sampling design for the collection of dove population data, using physiographic regions as the basis for stratification, was suggested by Foote, Peters, and Finkner (1958). The 79 regions designated in this report (fig. 1) are based essentially on a map entitled "Physical Divisions of the United States" prepared by Fenneman (1931). The boundaries of these divisions were modified in several instances after examination of field data and more recent ecological studies.




Figure 1.--Physiographic regions used in analysis of mourning dove population data, Revised 1970  
See page 5 for strata codes.

# Physiographic Regions Used in Analysis of Mourning Dove Population Data, Revised 1970 [Modified after Fenneman (1931)]

<u>Description</u>	<u>Stratum Code</u>	<u>Description</u>	<u>Stratum Code</u>	<u>Description</u>	<u>Stratum Code</u>
Laurentian Upland Division		Interior Plains Division		Intermontane Plateaus Division	
Superior Upland Province	010	Interior Low Plateaus Province		Columbia Plateaus Province	
Atlantic Plain Division		Highland Rim section	111	Walla Walla Plateau	201
Coastal Plain Province		Lexington Plain	112	Blue Mountain section	202
Embayed section	031	Nashville Basin	113	Payette section	203
Upper Coastal Plain	032	Central Lowland Province		Snake River Plain	204
Floridan section	033	Eastern lake section	121	Harney section	205
East Gulf Coastal Plain	034	Western lake section	122	Colorado Plateaus Province	
Mississippi Alluvial Plain	035	Wisconsin Driftless section	123	High Plateaus of Utah	211
West Gulf Coastal Plain	036	Till Plains	124	Uinta Basin	212
Lower Coastal Plain	037	Dissected Till Plains	125	Canyon Lands	213
Appalachian Highlands Division		Osage Plains	126	Navajo section	214
Piedmont Province		Great Plains Province		Grand Canyon section	215
Piedmont Uplands	041	Central Texas section	130	Datil section	216
Piedmont Lowlands	042	Missouri Plateau, glaciated	131	Basin and Range Province	
Blue Ridge Province		Missouri Plateau, unglaciated	132	Great Basin	221
Northern section	051	Black Hills	133	Sonoran Desert	222
Southern section	052	High Plains	134	Salton Trough	223
Valley and Ridge Province		Plains Border	135	Mexican Highland	224
Tennessee section	061	Colorado Piedmont	136	Sacramento section	225
Middle and Hudson Valley section	062	Raton section	137		
St. Lawrence Valley Province		Pecos Valley	138		
Champlain and Northern section		Edwards Plateau	139		
Appalachian Plateaus Province	070	Interior Highlands Division		Pacific Mountain Division	
Mohawk and Allegheny section	081	Ozark Plateaus Province		Cascade Sierra Mountains Province	
Catskill section	082	Springfield-Salem plateaus	141	Northern Cascade Mountains	231
Kanawha section	085	Boston "Mountains"	142	Middle Cascade Mountains	232
New England section	086	Ouachita Province		Southern Cascade Mountains	233
Southern New England section	091	Arkansas Valley	151	Sierra Nevada	234
Northern New England section	092	Ouachita Mountains	152	Pacific Border Province	
Mountain section	093	Rocky Mountain Division		Puget Trough	241
Taconic section	095	Southern Rocky Mountains Province	160	Olympic Mountains	242
Adirondack Province	100	Wyoming Basin Province	170	Oregon Coast Range	243
		Middle Rocky Mountains Province	180	Klamath Mountains	244
		Northern Rocky Mountains Province	190	California Trough	245
				California Coast Ranges	246
				Los Angeles Ranges	247
				Lower Californian Province	250

## Breeding Density Index

The Breeding Density Index (BDI) is an indicator of the number of doves per unit of area and is derived from the average number of calling doves per route. To obtain as accurate an average as possible for derivation of this index, the call-count data from each stratum in each State are weighted according to the land areas they represent.

Before 1966, the BDI for each State represented the average number of birds heard calling per route within that State, thus weighting all routes equally. The State averages were then weighted in proportion to the estimated area of dove habitat in each State of a management unit (fig. 2) to provide a Breeding Population Index for each unit (U.S. Bureau of Sport Fisheries and Wildlife, 1957).

Beginning with the 1966 survey, weighting factors based on physiographic regions were used for calculating BDI values in States which had been "randomized" for 2 or more years. The average number of doves heard calling per route in each region within a State was weighted by the percentage of the total land area in the State occupied by that region.

Calculation of management unit BDI's since 1965 has involved two similar procedures as a result of a computer program change. In 1966 and 1967, indexes were determined for each management unit by weighting each State's BDI by the percentage of the total land area occupied by that State in the management unit. When a region within a State was not represented by a BDI, that region assumed the mean of the other regions weighted by land area in that State. In 1968 and 1969, management unit BDI's were derived directly from State physiographic region BDI's. When a region within a State was not represented by a BDI, that region assumed the weighted management unit mean. Minor differences between these procedures are evident only when physiographic regions within States are not represented by comparable routes.

Random routes in Maine, New Hampshire, Rhode Island, and Vermont were established after initiation of a special study in 1966 to determine the effects of hunting regulations on dove populations in the Eastern Management Unit. In order to preserve the experimental design of the special study, call-count data from these States have not been included in the Eastern Management Unit means. In addition, no data were received from Delaware in 1969. Consequently, the 1969 and 1970 management unit means were derived from comparable routes run in 43 States.






Figure 2.--Mourning dove management units

### Determination of short-term population changes, 1969 to 1970

Changes in the size of mourning dove breeding populations between 1969 and 1970 are indicated by data from 679 comparable routes run in both years. Weighted average BDI values for each year are presented by States and management units. Differences in these BDI values, expressed as percent change, determined the magnitude of changes in the breeding population.

### Determination of long-term population trends by State, 1960 to 1970

Short-term (year-to-year) population changes are based, as indicated, on data from comparable routes only. Since the composition of these comparable routes changes with each 2-year comparison, a Base-Year (BY) has been chosen for each State. Long-term trends are shown by applying the percent change in the BDI from year to year to this index. Before 1967, the BY was generally the first year that the State's call-count routes were randomized. In order to provide a more uniform basis for evaluation of long-term trends, 1967 has been selected as the BY for all States. The index for this Base Year was obtained by taking the mean of comparable routes run in both 1966 and 1967 (Ruos and MacDonald, 1968). This BY value is thought to provide a meaningful refinement over the previous method. Not only are two "random-route" years averaged to reduce the influence of a possible atypical year, but the choice of a uniform BY for all States reduces possible bias in overweighting a State by the selection of a BY in a peak year.

As in the past, the BDI value for each State for each year is adjusted to the BY value and weighted to provide management unit values. This weighting is based upon differences in land area among States. The land area values and the adjusted BDI values for States and management units are presented.

### Determination of long-term trends by physiographic region, 1965 to 1970

Dove BDI's have been determined for each physiographic region since 1965. The 6-year trend has been obtained by adjusting these values to a base year similar to the procedures described in the preceding section, "Determination of long-term population trends by State, 1960 to 1970." 1969 has been selected as the Base-Year for all regions. In order to reduce the influence of low sampling intensity within regions, the 1969 BYI represents the average of the mean number of doves heard per route in 1968, 1969, and 1970.

The land area values and the BDI's adjusted to the 1969 BY are presented for each physiographic region within management units.

## Computer analysis of dove call-count data

Through the efforts of the North Carolina Institute of Statistics, University of North Carolina, and with the support of the Southeastern Association of Game and Fish Commissioners, an improved computer program was made available for the analysis of the 1968, 1969, and 1970 call-count data. This program provides properly weighted State and management unit averages. It yields the mean difference, the standard error of the mean difference, and the level of significance of the change for each State and management unit. This program also provides a summary of data by physiographic region irrespective of State boundaries, thus allowing analysis of population distribution by physiographic region.

## Statistical evaluation of data

The Call-Count Survey was designed to detect major year-to-year changes in the breeding population index within each management unit (Foote, 1959). Analysis of the 1969 and 1970 data revealed that observed differences of 8.5, 7.6, and 14.2 percent between these years within the Eastern, Central, and Western Management Units, respectively, would be statistically significant at the 95-percent level. For the entire country, an observed difference of 5.4 percent in the BDI between 1969 and 1970 would be significant. Although the survey was not designed to detect a change in the BDI between years within States or physiographic regions, data from these areas were also subjected to statistical analysis.

Long-term BDI's, adjusted to a BY for all physiographic regions, States, and management units, were examined to determine whether significant trends were present. Data from each source were analyzed using a linear regression model.

## Determination of population distribution

The density distribution of doves has been determined from a study of BDI values adjusted to a BY for each physiographic region and State. These data for 1970 have been assigned to one of five density classes. Changes in the adjusted BDI's greater than 10 percent between 1969 and 1970 within physiographic region and State also were determined.

## FINDINGS

From 1969 to 1970, the dove breeding population index increased by 4 percent in the Eastern Management Unit because of small increases in



the population indexes for the combined hunting States and the combined nonhunting States; but it decreased by 2 percent in the Central Unit because of a small increase for the combined hunting States and a large decrease for the combined nonhunting States; and by 11 percent in the Western Unit where all States are hunting States.

The Central and Western Management Unit population indexes are at their lowest levels for the 11-year period, 1960-70. All 1970 management unit values are below their preceding 10-year means. Further, a statistically significant downward population trend was found in each management unit. Additional study of these data is presented by management unit.

#### Status of the United States dove population

*1970 population distribution.*--The density distribution of mourning dove populations in the United States is presented by State (fig. 3) and by physiographic region (fig. 4). The most extensive area of high dove density was in the middle States, especially in the east-central Great Plains and Central Lowlands. Other important breeding population areas were observed in the Upper Coastal Plain of Georgia and the Carolinas and in the lower Mississippi River plain. A mean of 40 or more doves were heard per route in the States of Indiana, Kansas, and Nebraska.

*1969 to 1970 population changes.*--The United States BDI declined 2.1 percent from 19.1 doves heard per route in 1969 to 18.7 doves heard per route in 1970 (table 1). Changes greater than 10 percent in the BDI are illustrated by State (fig. 5) and by physiographic region (fig. 6). The 1970 indexes were generally lower than in 1969 in the northern and western areas of the country, and generally higher in sections of the Rocky Mountains, mid-States, and eastern seaboard. From 1969 to 1970, no change occurred in the BDI of 19.7 doves heard per route for the combined hunting States; whereas a decline of 8.1 percent from 17.3 to 15.9 doves heard per route occurred in the combined nonhunting States index.

*1960 to 1970 long-term population trends.*--The 1970 BDI's adjusted to a Base-Year for the United States, the combined hunting States, and the combined nonhunting States are the lowest observed for the 11-year period, 1960-1970 (table 2). This is the fourth successive year without an increase in adjusted BDI's. The adjusted BDI's for these areas in 1970 are also well below the preceding 10-year means: United States, -14.2 percent; hunting States, -11.9 percent; and nonhunting States, -19.9 percent.




Figure 3.--Relative densities of breeding mourning doves by State, 1970






Figure 4.--Relative densities of breeding mourning doves by physiographic region, 1970




Figure 5.--Changes in densities of breeding mourning doves by State between 1969 and 1970






Figure 6.--Changes in densities of breeding mourning doves by physiographic region between 1969 and 1970

Adjusted BDI's plotted in figures 7 and 8 reflect the general downward trend in population indexes since 1960. Linear regression analyses of these data (table 2) are shown in figure 9. The adjusted BDI's declined at an average annual rate of 2.8 percent in the United States, 2.5 percent in the hunting States, and 3.3 percent in the non-hunting States. Thus, this study reveals a significant decline in U.S. populations between 1960 and 1970.

Population trends as determined from linear regression analyses are shown by State (table 2, fig. 10) and by physiographic region (table 3, fig. 11). From 1960 to 1970, statistically significant downward trends exist in much of the northern Great Plains and the Pacific border and southern border States. Trends are significantly upward in several scattered mid-latitude States.

#### Status of the Eastern Management Unit population

*1970 population distribution.*--Highest dove population densities in the Eastern Management Unit were in the west-central section, especially in the Central Lowlands. Densities were generally low in the Appalachian Highlands, northern uplands, and portions of the lower Atlantic Coastal Plain (fig. 4). States represented by a mean of 30 or more doves heard per route included Indiana, Ohio, Tennessee, Mississippi, and New Jersey (fig. 3).

*1969 to 1970 population changes.*--The Eastern Management Unit BDI increased 3.8 percent from 18.3 doves heard per route in 1969 to 19.0 doves heard per route in 1970 (table 1). The 1970 population levels were generally higher than in 1969 along the Atlantic seaboard and in sections of the Central Lowland and lower Mississippi plain. Population indexes were lower in the northern border States (figs. 5, 6). From 1969 to 1970, the combined hunting States index increased 4.1 percent and the combined nonhunting States index increased 1.9 percent.

*1960 to 1970 long-term population trends.*--The increase in the adjusted unit-wide BDI from 1969 to 1970 followed 3 successive years of annual population decrease (table 2). Population levels increased for the first time in 5 years in the combined hunting States, while the combined nonhunting States index increased for the second successive year. The adjusted unit BDI for 1970 is 3.1 percent below the preceding 10-year mean (fig. 7). The combined hunting States 1970 index is 7.2 percent below this mean. In contrast, the adjusted 1970 BDI for the combined nonhunting States is 8.0 percent above the preceding 10-year mean (fig. 12).




Figure 7.--Breeding mourning dove population indexes by management unit, 1960-1970






Figure 8.--Breeding mourning dove population indexes for the United States hunting and nonhunting States, 1960-1970




Figure 9.--Linear regression lines of mourning dove call-count data, 1960-1970




Figure 10.--Trends in mourning dove breeding populations by State, 1960 to 1970








Figure 12.--Breeding mourning dove population indexes for the Eastern and Central Management Unit hunting and nonhunting States, 1960-1970

Regression analysis shows a significant downward trend in the Eastern Unit population between 1960 and 1970; the mean rate of decline was determined to be 1.0 percent per year (table 2). During the same period, the combined hunting States index declined significantly at an average annual rate of 1.8 percent. Although no significant trend was found for the nonhunting States, the index increased at an average rate of 1.3 percent per year (fig. 9). Significant upward population trends were limited to Indiana, Ohio, Maryland, and South Carolina, whereas downward trends were most common in the northern and southern border States (figs. 10, 11).

#### Status of the Central Management Unit population

*1970 population distribution.*--Highest population densities in the Central Management Unit were distributed in the east-central section, especially in the Great Plains and Central Lowlands. Low densities were rather uniformly distributed in the northern, western, and southern parts of the Unit (fig. 4). States represented by a mean of 30 or more doves heard per route included Kansas, Nebraska, South Dakota, Oklahoma, and Missouri (fig. 3).

*1969 to 1970 population changes.*--The Central Management Unit BDI declined 1.8 percent from 22.1 doves heard per route in 1969 to 21.7 doves heard per route in 1970 (table 1). The 1970 population levels remained the same or were generally higher than in 1969 in the southeastern part of the Unit. Population indexes were lower in the northern and western sections (figs. 5, 6). From 1969 to 1970, the combined hunting States index increased 3.3 percent, whereas the combined nonhunting States index decreased 12.8 percent.

*1960 to 1970 long-term population trends.*-- In 1970, the Central Unit BDI adjusted to a Base-Year declined to its lowest level for the 11-year period (table 2). This represents the fourth successive annual decline. The population index also declined for the fourth successive year in the combined nonhunting States, while the population index for the combined hunting States increased from its near record low in 1969. Current population levels are well below the preceding 10-year means: Central Unit, -14.8 percent; combined hunting States, -6.4 percent; and combined nonhunting States, -29.6 percent (figs. 7, 12).

Regression analysis shows that a significant downward trend in dove population indexes occurred from 1960 to 1970 in the Central Unit. Similar downward trends were observed for the combined hunting States and combined nonhunting States (table 2). The annual rates of decline in the adjusted BDI's were determined as follows: Central Unit, -3.1

percent; combined hunting States, -2.2 percent; and combined nonhunting States, -4.9 percent (fig. 9). Significant upward population trends were limited to two States, Arkansas and Wyoming. Population trends were significantly downward in States bordering the Missouri River and in southern border States (figs. 10, 11).

#### Status of the Western Management Unit population

*1970 population distribution.*--Highest population densities in the Western Management Unit were essentially restricted to coastal California and sections of the Columbia Plateau. Lowest densities were distributed through much of the Great Basin and the northwestern part of the Western Unit (fig. 4). Idaho had the highest adjusted BDI of any Western Unit State in 1970, with 22.3 doves heard calling per route (table 2; fig. 3).

*1969 to 1970 population changes.*--The Western Management Unit BDI declined 11.2 percent from 14.3 doves heard per route in 1969 to 12.7 doves heard per route in 1970 (table 1). The 1970 population levels were generally lower throughout the Unit. Populations increased in several regions, especially in the States of Idaho and Washington (figs. 5, 6). All States within the Western Unit hunt mourning doves.

*1960 to 1970 population trends.*--The 1970 BDI adjusted to a Base-Year for the Western Unit is the lowest observed for the 11-year period (table 2). This is the sixth successive year without an increase in the BDI. The population index for this Unit in 1970 is 27.1 percent below the preceding 10-year mean (fig. 7).

Regression analysis shows a significant downward trend in the dove population index from 1960 to 1970. The adjusted BDI's declined at an average annual rate of 4.0 percent along the calculated regression line (fig. 9). Between 1960 and 1970, only the State of Nevada had a significant upward population trend. Significant downward trends occurred in California and Oregon during the same period (figs. 10, 11).

#### Statistical significance of data

*1969 to 1970 population changes.*--No significant ( $p < 0.05$ ) changes occurred in the BDI's of any management unit or in the combined hunting or nonhunting States of any unit between 1969 and 1970. Although not designed to detect population changes within States, the survey showed significant increases in Tennessee and Missouri. In no State did the index decrease significantly (table 1).

A study of physiographic region data obtained from the 1969 and 1970 call-count surveys was also made. As identified in figure 1, significant



( $p < 0.05$ ) increases occurred in sections of the Atlantic Coastal Plain (Regions 031 and 035), the Valley and Ridge Province of the Appalachian Highlands (061), and in the Salton Trough of southern California (223). Significant decreases occurred in the Sonoran-Mojave Desert (222) and in part of the Interior Highlands of Arkansas and Oklahoma (151).

*1960 to 1970 long-term population trends*.--Statistical analyses of the 1960-70 data revealed that significant ( $p < 0.05$ ) downward trends in adjusted BDI's occurred in all management units and submanagement units, except in the combined nonhunting States of the Eastern Unit (table 2). No statistical significance could be attached to the observed upward trend shown for the nonhunting States of the Eastern Unit.

For the first time, this report presents the results of long-term BDI data analyzed by State and physiographic region (tables 2 and 3). Seven States were determined to have significant ( $p > .95$ ) upward population trends between 1960 and 1970, while 16 States had downward trends in population (fig. 10). From 1965 to 1970, four of 71 physiographic regions had significant upward population trends, whereas 11 regions had downward trends in population (fig. 11).

#### ACKNOWLEDGEMENTS

This report would not be possible without the cooperation of the State conservation departments and the many individuals who conscientiously assisted in collecting data. Preparation of this report represents a combined effort; special acknowledgement is made to Mrs. Jaynie Peters for electronic data processing assistance, Dr. John P. Rogers for editing, and Mrs. Katheryn Munson and Mrs. Kathy Judy for typing the final manuscript.

## REFERENCES

Fenneman, Nevin M.

1931. Physiography of western United States. McGraw-Hill Book Company, New York. 534 p.

---

1938. Physiography of eastern United States. McGraw-Hill Book Company, New York. 714 p.

Foote, Leonard E.

1959. A sampling design for mourning dove call counts. A report to the Bureau of Sport Fisheries and Wildlife, U.S. Fish and Wildlife Service, from The Wildlife Management Institute. (Unpublished report.) December. 47 p. + 24 tables + appendix.

---

, and Harold S. Peters.

1952. Introduction, p. 1-3 in Investigations of methods of appraising the abundance of mourning doves. U.S. Fish and Wildlife Service, Special Scientific Report--Wildlife No. 17.

---

, Harold S. Peters, and Alva L. Finkner.

1958. Design tests for mourning dove call-count sampling in seven southeastern States. Journal of Wildlife Management, vol. 22, No. 4, p. 402-403.

Frankel, Arthur I., and Thomas S. Baskett.

1961. The effect of pairing on cooing of penned mourning doves. Journal of Wildlife Management, vol. 25, No. 4, p. 372-384.

Irby, Harold D.

1964. The relationship of calling behavior to mourning dove populations and production in southern Arizona. Unpublished Ph.D. dissertation, University of Arizona. 100 p.

Jackson, Gary L., and Thomas S. Baskett.

1964. Perch-cooing and other aspects of breeding behavior of mourning doves. Journal of Wildlife Management, vol. 28, No. 2, p. 293-307.

Ruos, James L., and Duncan MacDonald.

1968. Mourning dove status report, 1967. U.S. Bureau of Sport Fisheries and Wildlife, Special Scientific Report--Wildlife No. 121. 23 p.

---

, and Duncan MacDonald.

1970. Mourning dove status report, 1968. U.S. Bureau of Sport Fisheries and Wildlife, Special Scientific Report--Wildlife No. 129. 38 p.

Ruos, James L.

1970. Mourning dove status report, 1969. U.S. Bureau of Sport Fisheries and Wildlife, Special Scientific Report--Wildlife No. 132. 35 p.

(U.S.) Bureau of Sport Fisheries and Wildlife.

1957. Mourning Dove Newsletter, No. 12. 30 p.

Wight, Howard M.

1964. Matedness in the mourning dove and its effect on the nationwide dove-call census. Trans. 29th North American Wildlife and Resources Conference, p. 270-281.

## TABLES

Table 1.--Changes in mourning dove breeding density indexes, 1969-70

State	Comparable routes	Average doves heard/route (weighted) <u>1/</u>		Percent change <u>2/</u>
		1969	1970	
<u>EASTERN MANAGEMENT UNIT - HUNTING STATES</u>				
Ala.	28	21.3	18.7	- 12.2
Del.	0			
Fla.	20	7.4	8.5	+ 14.9
Ga.	21	18.2	22.2	+ 22.0
Ill.	13	23.4	28.0	+ 19.7
Ky.	11	37.3	34.5	- 7.5
La.	19	10.2	10.2	0.0
Md.	8	18.1	21.6	+ 19.3
Miss.	19	25.8	27.2	+ 5.4
N.C.	18	29.7	30.5	+ 2.7
Pa.	15	12.4	9.0	- 27.4
R.I. <u>3/</u>	2	13.0	4.5	- 65.4
S.C.	17	28.0	23.1	- 17.5
Tenn.	20	17.4	26.9	+ 54.6**
Va.	8	22.1	19.6	- 11.3
W. Va.	7	4.4	5.1	+ 15.9
Subtotal	226	19.4	20.2	+ 4.1
<u>EASTERN MANAGEMENT UNIT - NONHUNTING STATES</u>				
Conn.	2	1.5	5.5	+266.7
Ind.	11	34.7	35.5	+ 2.3
Maine <u>3/</u>	3	1.0	0.0	-100.0
Mass.	2	1.9	8.1	+326.3
Mich.	17	9.7	8.7	- 10.3
N.H. <u>3/</u>	4	3.0	3.2	+ 6.7
N.J.	3	20.3	25.4	+ 25.1
N.Y.	11	7.1	5.7	- 19.7
Ohio	10	28.3	31.4	+ 11.0
Vt. <u>3/</u>	3	0.1	0.4	+300.0
Wis.	15	10.4	9.3	- 10.6
Subtotal	81	16.0	16.3	+ 1.9
Eastern Unit				
Total	307	18.3	19.0	+ 3.8

See footnotes at end of table, p. 29.

Table 1.--Changes in mourning dove breeding density indexes, 1969-70--  
continued

State	Comparable routes	Average doves heard/route (weighted) <u>1/</u>		Percent change <u>2/</u>
		1969	1970	
<u>CENTRAL MANAGEMENT UNIT - HUNTING STATES</u>				
Ark.	12	23.6	24.3	+ 3.0
Colo.	9	13.5	14.1	+ 4.4
Kans.	16	58.8	54.4	- 7.5
Mo.	19	23.9	29.5	+23.4**
N. Mex.	14	15.6	14.7	- 5.8
Okla.	9	32.5	29.6	- 8.9
S. Dak. <sup>4/</sup>	10	32.7	34.7	+ 6.1
Tex.	43	17.6	20.1	+14.2
Sub total	132	24.6	25.4	+ 3.3
<u>CENTRAL MANAGEMENT UNIT - NONHUNTING STATES</u>				
Iowa	12	25.8	19.1	-26.0
Minn.	10	8.3	6.8	-18.1*
Mont.	9	9.8	7.4	-24.5
Nebr.	21	41.4	40.9	- 1.2
N. Dak.	20	22.3	18.9	-15.2
Wyo.	9	8.8	7.8	-11.4
Sub total	81	17.9	15.6	-12.8*
Central Unit Total	213	22.1	21.7	- 1.8

See footnotes at end of table, p. 29.

Table 1.--Changes in mourning dove breeding density indexes, 1969-70--  
continued

State	Comparable routes	Average doves heard/route (weighted) <u>1/</u>		Percent change <u>2/</u>
		1969	1970	
<u>WESTERN MANAGEMENT UNIT</u>				
Ariz.	40	31.7	23.0	-27.4*
Calif.	49	12.1	11.5	- 5.0
Idaho	11	13.9	18.2	+30.9
Nev.	12	8.2	7.5	- 8.5
Oreg.	18	10.8	7.4	-31.5
Utah	12	10.3	8.5	-17.5
Wash.	17	10.3	11.4	+10.7
<hr/>				
Western Unit Total	159	14.3	12.7	-11.2
<hr/>				
U.S. Hunt	517	19.7	19.7	± 0
U.S. Nonhunt	162	17.3	15.9	- 8.1
U.S. Total	679	19.1	18.7	- 2.1

1/ Except as noted, State and management unit indexes were obtained from comparable, randomized route data adjusted for variation in the land area of each physiographic region represented.

2/ Probability that observed change represents actual change: \*90 percent; \*\*95 percent; \*\*\*99 percent.

3/ Rhode Island and Vermont data from randomized routes, Maine and New Hampshire from management routes. These data not represented in management unit means to preserve comparability for Eastern Management Unit dove study.

4/ South Dakota included as nonhunting State in previous reports. Dove hunting reestablished in South Dakota since 1967-68 season.



Table 2.--Trends in mourning dove breeding density indexes by State, 1960-70

State	Weight factor <sup>1/</sup>	ADJUSTED AVERAGE DOVES HEARD CALLING PER ROUTE $\frac{2}{1960}$ $\frac{2}{1961}$ $\frac{2}{1962}$ $\frac{2}{1963}$ $\frac{2}{1964}$ $\frac{2}{1965}$ $\frac{2}{1966}$ $\frac{2}{1967}$ $\frac{2}{1968}$ $\frac{2}{1969}$ $\frac{2}{1970}$											LINEAR REGRESSION, 1960-1970		Stat. Sign. of Trend
													Percent	No. of Doves	
EASTERN MANAGEMENT UNIT - HUNTING STATES															
Ala.	33.32	22.0	19.4	18.9	16.3	21.1	20.5	20.5	19.8	17.1	20.8	18.3	- 0.6	-0.1	n.s. <sup>4/</sup>
Del.	1.29	6.2	4.6	5.6	5.1	4.4	5.1	3.8	7.5	5.2	--	--	--	--	--
Fla.	35.82	10.1	9.4	11.4	9.8	11.2	10.5	9.4	9.1	7.6	7.8	9.0	- 2.5	-0.2	p .95
Ga.	37.82	16.9	14.4	16.8	14.4	22.5	18.5	13.1	15.4	14.2	15.5	18.9	0.0	0.0	n.s.
Ill.	35.09	28.3	32.3	26.3	26.9	24.3	23.5	27.0	27.9	27.0	22.1	26.5	- 1.5	-0.4	n.s.
Ky.	26.08	29.8	28.2	28.6	27.0	30.4	32.5	35.3	29.9	26.2	26.1	24.2	- 1.0	-0.3	n.s.
La.	31.14	19.3	17.8	14.2	16.6	13.6	11.6	7.7	8.9	7.4	7.5	7.5	-11.2	-1.3	p .99
Md.	6.55	15.6	14.4	15.8	16.1	16.1	16.0	15.2	20.1	17.5	16.7	20.0	+ 2.4	+0.4	p .95
Miss.	30.63	37.8	33.2	35.4	30.5	35.1	36.8	33.6	27.7	26.4	28.5	30.1	- 2.6	-0.8	p .95
N.C.	22.51	19.7	17.3	19.0	21.0	21.9	24.8	26.4	21.4	22.7	18.4	18.9	+ 0.7	+0.2	n.s.
Pa.	29.01	9.5	9.9	8.2	5.6	7.2	7.4	8.7	11.9	9.0	9.8	7.1	+ 0.4	0.0	n.s.
R.I. <sup>5/</sup>	0.67	4.0	5.0	5.5	5.5	8.9	5.9	4.2	5.3	8.8	8.2	2.9	+ 1.9	+0.1	n.s.
S.C.	19.99	25.8	25.5	23.0	23.0	24.1	35.3	34.4	34.2	33.1	33.8	27.9	+ 3.4	+1.0	p .95
Tenn.	27.07	35.6	30.2	34.8	29.2	28.1	30.4	30.4	22.3	23.5	22.5	34.8	- 2.5	-0.7	n.s.
Va.	26.05	30.6	28.7	24.8	25.2	29.0	24.0	29.1	23.6	31.8	24.4	21.6	- 1.5	-0.4	n.s.
W. Va.	15.41	20.1	21.0	44.0	18.5	23.4	22.6	9.8	5.0	5.0	5.7	6.6	-18.5	-2.6	p .95
SUB-TOT/															
AVG.	377.78	23.0	21.5	22.2	19.8	21.9	22.0	21.2	19.6	18.9	18.4	19.3	- 1.8	-0.4	p .99
EASTERN MANAGEMENT UNIT - NONHUNTING STATES															
Conn.	3.23	6.5	3.5	3.5	3.0	2.4	1.9	2.9	4.5	4.9	1.2	4.5	- 2.9	-0.1	n.s.
Ind.	23.36	25.1	28.3	23.4	26.8	27.7	20.2	37.2	38.6	41.7	40.2	41.1	+ 6.6	+2.0	p .99
Maine <sup>5/</sup>	19.85	3.0	3.0	3.0	1.3	0.0	0.3	1.3	0.0	1.0	1.5	0.0	-28.0	-0.3	p .95
Mass.	5.31	10.6	6.9	9.0	5.2	6.7	7.7	11.0	14.5	4.9	1.4	6.1	- 4.0	-0.3	n.s.
Mich.	37.18	14.0	14.8	11.8	17.4	13.2	8.0	12.3	11.9	9.1	11.8	10.6	- 3.6	-0.4	p .90
N.H. <sup>5/</sup>	5.80	6.8	4.6	3.4	3.1	6.3	10.6	7.5	5.0	4.4	4.0	4.3	- 1.1	-0.1	n.s.
N.N.J.	4.91	32.7	29.0	28.8	29.5	30.4	25.7	24.2	20.1	30.0	26.3	32.9	- 1.0	-0.3	n.s.
N.Y.	30.49	8.8	7.2	8.3	7.8	7.0	7.1	7.9	7.8	6.8	5.8	4.7	- 3.7	-0.3	p .99
Ohio	26.42	14.8	17.9	21.4	17.5	16.8	19.1	26.3	25.5	27.0	30.6	34.0	+ 8.3	+1.7	p .99
Vt. <sup>5/</sup>	5.95	8.0	0.0	2.0	2.0	2.0	1.0	6.0	5.0	3.5	2.0	8.0	+ 6.0	+0.2	n.s.
Wis.	36.07	18.3	18.1	12.7	14.1	15.2	16.1	11.1	13.5	11.4	9.8	8.8	- 6.0	-0.8	p .99
SUB-TOT/															
AVG.	166.97	16.0	16.4	14.7	16.0	15.2	13.4	17.1	17.6	17.0	17.1	17.5	+ 1.3	+0.2	n.s.
EMU TOT/															
AVG.	544.75	20.8	20.0	19.9	18.6	19.8	19.4	20.0	19.0	18.3	18.0	18.8	- 1.0	-0.2	p .99

See footnotes at end of table, p. 32.



Table 2--Trends in mourning dove breeding density indexes by State, 1960-1970--continued

State	Weight factor <sup>1/</sup>	ADJUSTED AVERAGE DOVES HEARD CALLING PER ROUTE $\frac{2}{1960}$ $\frac{3}{1970}$										LINEAR REGRESSION, 1960-1970			
												Mean Annual Chg. (BYI)	Stat. Sign.		
												Percent	of Trend		
CENTRAL MANAGEMENT UNIT - HUNTING STATES															
Ark.	34.37	17.4	19.0	10.5	16.0	16.8	14.8	16.5	20.7	20.0	22.3	22.9	+4.1	+0.7	p .95
Colo.	67.18	28.8	13.8	13.8	11.0	13.1	11.5	14.8	16.1	13.7	19.7	20.5	-0.4	-0.1	n.s.
Kans.	52.43	54.4	55.2	56.2	49.8	53.1	48.2	51.8	60.3	59.2	65.6	60.7	+1.7	+0.9	p .90
Mo.	45.10	50.2	45.2	42.7	36.7	40.2	36.0	39.3	36.6	39.1	25.2	31.1	-4.4	-1.7	p .99
N. Mex.	77.98	32.5	21.8	19.2	18.1	20.6	19.3	28.4	8.5	16.0	11.0	10.4	-8.6	-1.6	p .95
Okla.	44.40	40.9	41.6	36.4	39.9	37.7	34.6	29.3	37.9	41.3	33.9	30.9	-1.9	-0.7	p .90
S. Dak.	49.20	40.2	48.5	37.4	40.6	39.0	35.1	45.3	28.2	32.9	30.1	31.9	-3.8	-1.3	p .95
Tex.	170.03	22.9	18.6	19.4	17.7	17.4	16.4	17.8	16.8	15.8	15.1	17.3	-2.8	-0.5	p .99
SUB-TOT/ AVG.															
540.69		33.0	28.9	26.7	25.4	26.3	24.0	27.4	24.4	25.7	24.2	24.9	-2.2	-0.6	p .95
CENTRAL MANAGEMENT UNIT - NONHUNTING STATES															
Iowa	36.15	30.6	38.3	37.1	33.5	34.9	29.0	33.2	34.1	30.7	25.8	19.1	-3.6	-1.2	p .95
Minn.	54.09	15.0	19.3	15.9	16.2	20.5	18.6	18.7	16.7	19.1	10.0	8.2	-3.5	-0.6	n.s.
Mont.	94.47	18.3	15.7	10.4	15.1	15.4	15.7	17.1	18.7	5.7	6.6	5.0	-7.6	-1.0	p .95
Nebr.	49.69	109.3	88.6	69.9	58.7	66.1	55.6	48.5	40.0	47.8	47.7	47.2	-8.9	-5.4	p .99
N. Dak.	45.54	21.2	22.3	20.1	20.0	20.2	23.4	20.7	20.7	24.2	20.2	17.2	-0.6	-0.1	n.s.
Wyo.	62.33	10.0	11.8	10.8	12.7	9.8	12.1	15.9	13.0	9.3	20.6	18.3	+5.8	+0.7	p .95
SUB-TOT/ AVG.															
342.27		31.2	29.4	24.1	23.8	25.2	23.7	23.9	22.3	19.7	19.5	17.1	-4.9	-1.2	p .99
CMU TOT/ AVG.															
882.96		32.3	29.1	25.7	24.8	25.9	23.9	26.1	23.6	23.3	22.4	21.9	-3.1	-0.8	p .99

See footnotes at end of table, p. 32.

Table 2.--Trends in mourning dove breeding density indexes by State, 1960-1970--continued

State	Weight factor <sup>1/</sup>	ADJUSTED AVERAGE DOVES HEARD CALLING PER ROUTE 2/ 3/ 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970											LINEAR REGRESSION, 1960-1970		
													Mean Annual Chg.(BYI)	Stat.Sign.	
													Percent	of Trend	
WESTERN MANAGEMENT UNIT															
Ariz.	72.65	19.1	34.3	23.8	22.7	27.6	26.6	30.4	27.6	23.8	27.0	19.6	- 0.5	-0.1	n.s.
Calif.	101.71	36.2	26.0	29.2	29.2	32.0	21.6	17.3	14.2	12.0	11.5	11.0	-12.4	-2.5	p .99
Idaho	54.37	20.4	18.0	18.3	18.3	20.6	19.4	18.6	17.8	18.0	17.0	22.3	+ 0.1	0.0	n.s.
Nev.	71.27	1.9	2.3	1.2	1.8	2.9	2.7	4.0	4.4	8.2	6.3	5.8	+22.7	+0.6	p .99
Oreg.	62.27	19.2	21.2	18.2	16.8	18.1	14.1	14.0	12.9	13.2	14.9	10.2	- 5.6	-0.9	p .99
Utah	53.34	18.6	14.8	13.6	13.4	12.9	13.7	14.8	22.2	12.3	8.6	7.1	- 4.4	-0.6	n.s.
Wash.	43.87	10.5	9.2	11.0	9.8	13.2	13.1	13.3	11.4	10.5	11.1	12.3	+ 1.5	+0.2	n.s.
WMU TOT/ AVG.	459.48	19.5	19.1	17.7	17.3	19.5	16.4	16.3	15.7	14.0	13.9	12.4	- 4.0	-0.7	p .99
TOTAL HUNTING STATES	1,377.95	25.8	23.6	22.4	21.2	22.8	20.9	22.0	20.2	19.9	19.2	19.2	- 2.5	-0.6	p .99
TOTAL NONHUNT. STATES	509.24	26.2	25.2	21.0	21.2	22.4	20.4	21.6	20.8	18.8	18.7	17.3	- 3.3	-0.7	p .99
U.S.TOT/ AVG.	1,887.19	25.9	24.0	22.0	21.2	22.7	20.8	21.9	20.4	19.6	19.0	18.7	- 2.8	-0.6	p .99

<sup>1/</sup> Assigned State land-area value.

<sup>2/</sup> From a Base Year, average doves heard calling per route for each State has been adjusted annually according to the percent change from the preceding year in counts on comparable routes. Except as noted, percent changes between years have been based upon unweighted, randomized data for the period 1959 to 1965 inclusive. Annual percent changes since 1966 have been derived from randomized data weighted by physiographic regions within States. The Base Year (mean of 1966 and 1967) was changed from previous reports in 1967 (Ruos and MacDonald, 1968).

<sup>3/</sup> Percent departures calculated from data carried to three decimal places; hence, apparent rounding errors.

Unit and subunit averages calculated by weighting individual State indexes; hence, change in unit long-term-trend index may not indicate precisely the same percent change as the 2-year comparison.

<sup>4/</sup> Not statistically significant ( $p < .90$ ).

<sup>5/</sup> Indexes for Maine, New Hampshire, Rhode Island, and Vermont have not been adjusted to a random base-year and have not been included in the unit "weighted means."

<sup>6/</sup> South Dakota included as a nonhunting State in previous reports. Hunting permitted in 1967, 1968, 1969.

Table 3.--Trends in mourning dove breeding density indexes by physiographic region, 1965-70

Region	Weight <sub>1</sub> / factor	ADJUSTED AVERAGE DOVES HEARD/ROUTE $\frac{2}{3}/$						LINEAR REGRESSION, 1965-1970			
								Mean Annual Chg. (BYI) Stat. Sign.			
		1965	1966	1967	1968	1969	1970	Percent	No. of Doves	of Trend	
EASTERN MANAGEMENT UNIT											
010	32.10	8.8	4.3	3.8	3.6	2.8	2.0	-26.3	-1.1	p .95	
031	15.34	29.5	30.3	28.5	28.2	21.5	24.9	- 5.1	-1.4	p .90	
032	16.40	53.0	39.3	45.8	47.9	44.5	39.6	- 3.1	-1.4	n.s.	
033	24.79	13.4	10.2	10.1	9.1	9.1	11.5	- 3.7	-0.4	n.s.	
034	63.72	26.8	22.9	20.6	17.7	20.4	21.2	- 5.0	-1.1	n.s.	
035	20.50	27.1	28.4	29.1	27.4	28.7	34.8	+ 3.8	+1.1	n.s.	
036	15.69	18.1	19.6	18.0	16.6	16.3	16.3	- 3.3	-0.6	p .95	
037	33.14	12.1	12.7	11.8	10.5	16.0	17.6	+ 8.1	+1.0	n.s.	
041	39.35	19.3	21.1	17.6	22.8	17.9	16.7	- 2.6	-0.5	n.s.	
042	3.51	14.6	24.1	24.6	35.7	31.4	27.3	+11.4	+2.8	n.s.	
051	1.93	No Estimate						n.e.	n.e.	n.e.	
052	6.09	10.0	13.2	11.2	13.2	7.7	11.8	- 1.4	-0.2	n.s.	
061	17.62	25.6	25.6	20.5	16.4	19.1	33.0	+ 1.7	+0.4	n.s.	
062	18.99	17.0	15.4	12.2	15.6	16.8	14.4	- 1.0	-0.2	n.s.	
070	2.40	No Estimate						n.e.	n.e.	n.e.	
081	27.02	7.4	9.2	14.5	12.2	9.3	7.9	+ 0.1	0.0	n.s.	
082	1.32	n.e.	3.0	0.0	1.0	0.0	0.0	-62.0	-0.6	n.s.	
085	32.72	5.4	6.5	4.9	3.3	4.1	3.7	-10.3	-0.5	p .90	
086	7.27	11.2	10.8	7.9	9.4	9.7	9.5	- 3.0	-0.3	n.s.	
091	9.28	n.e.	29.7	40.4	19.7	5.2	14.9	-28.2	-6.5	n.s.	
092	10.00	No	Estimate						n.e.	n.e.	n.e.
093	20.12	No	Estimate						n.e.	n.e.	n.e.
095	1.64	n.e.	n.e.	1.4	1.4	1.4	3.0	n.e.	n.e.	n.e.	
100	6.71	n.e.	0.7	0.0	0.0	0.0	0.0	-70.8	-0.1	n.s.	
111	25.51	43.3	45.1	39.3	38.5	35.6	35.1	- 5.0	-2.0	p .99	
112	6.70	18.2	31.3	17.0	14.2	20.8	17.7	- 5.2	-1.1	n.s.	
113	2.07	27.0	27.0	10.9	17.6	15.2	17.5	-11.0	-2.2	n.s.	
121	46.46	10.0	15.6	15.7	12.6	15.3	13.1	+ 2.4	+0.3	n.s.	
123	12.09	22.7	18.6	23.0	19.8	15.3	16.0	- 6.8	-1.3	p .90	
124	56.27	28.2	32.9	35.3	36.3	34.1	40.1	+ 5.5	+1.8	p .95	
141	0.27	30.3	26.2	32.6	33.5	27.8	30.7	+ 0.7	+0.2	n.s.	
EMU TOT/ AVG. 4/	577.02	19.4	20.0	19.0	18.3	18.0	18.8	- 1.0	-0.2	p .99	

See footnotes at end of table, p.36.

Table 3.--Trends in mourning dove breeding density indexes by physiographic region,  
1965-1970--continued

Region	Weight <sub>1</sub> / factor	ADJUSTED AVERAGE DOVES HEARD/ROUTE					2/3/ 1969 1970		LINEAR REGRESSION, 1965-1970			
		1965	1966	1967	1968	1969	1970	Mean Annual Chg. (BYI)	Percent No. of Doves	Stat. Sign. of Trend		
CENTRAL MANAGEMENT UNIT												
010	30.37	8.8	4.3	3.8	3.6	2.8	2.0	-26.3	- 1.1	p .95		
035	16.60	27.1	28.4	29.1	27.4	28.7	34.8	+ 3.8	+ 1.1	n.s.		
036	77.45	18.1	19.6	18.0	16.6	16.3	16.3	- 3.3	- 0.6	p .95		
122	64.68	34.1	37.0	33.6	35.5	27.4	25.1	- 6.2	- 2.1	p .90		
123	2.54	22.7	18.6	23.0	19.8	15.3	16.0	- 6.8	- 1.3	p .90		
124	0.82	28.2	32.9	35.3	36.3	34.1	40.1	+ 5.5	+ 1.8	p .95		
125	58.68	38.9	46.5	43.1	50.8	37.7	33.2	- 3.2	- 1.3	n.s.		
126	71.40	35.4	36.0	43.3	43.6	37.9	43.1	+ 3.2	+ 1.3	n.s.		
130	14.35	50.9	67.8	37.4	49.3	45.3	41.5	- 5.9	- 2.9	n.s.		
131	46.47	18.5	15.7	12.9	20.7	24.1	19.6	+ 6.1	+ 1.1	n.s.		
132	81.54	41.1	45.0	37.1	23.0	21.0	18.4	-17.9	- 5.7	p .99		
133	2.89	5.0	5.0	0.0	n.e.	41.0	71.1	n.s.	n.s.	n.s.		
134	97.09	27.5	22.0	20.0	24.2	23.8	22.9	- 1.6	- 0.4	n.s.		
135	25.78	50.8	53.1	61.7	59.3	64.8	66.7	+ 5.6	+ 3.2	p .99		
136	16.40	16.6	15.4	14.7	12.6	19.4	20.7	+ 5.4	+ 0.9	n.s.		
137	10.54	2.3	11.0	4.2	6.1	7.4	11.1	+16.2	+ 1.5	n.s.		
138	19.75	114.3	165.0	28.7	41.6	14.8	17.3	-46.5	-26.4	p .90		
139	22.05	1.8	1.6	24.5	8.7	19.8	17.0	+40.0	+ 3.3	n.s.		
141	27.75	30.3	26.2	32.6	33.5	27.8	30.7	+ 0.7	+ 0.2	n.s.		
142	3.11	19.4	7.4	5.0	2.3	3.5	2.3	-47.4	- 2.9	p .95		
151	4.70	6.7	20.2	13.0	21.0	21.5	15.9	+10.9	+ 1.7	n.s.		
152	6.98	3.0	3.0	3.3	3.9	4.9	3.3	+ 6.5	+ 0.2	n.s.		
160	28.53	2.9	5.1	2.6	4.4	7.5	5.2	+14.1	+ 0.6	n.s.		
170	24.23	10.9	14.5	17.2	8.1	13.4	17.9	+ 4.9	+ 0.6	n.s.		
180	23.58	3.7	5.6	6.4	7.7	7.7	7.4	+12.7	+ 0.7	p .95		
190	33.18	8.2	9.9	7.1	6.1	7.2	8.2	- 3.3	- 0.3	n.s.		
212	4.68	8.2	22.7	55.7	9.9	14.4	5.4	-12.1	- 2.4	n.s.		
213	6.02	0.9	1.4	4.0	4.2	6.2	8.1	+89.5	+ 1.4	p .99		
214	11.49	8.3	16.9	7.1	9.0	10.3	8.8	- 4.3	- 0.5	n.s.		
216	7.07	84.7	75.6	28.7	46.5	20.6	17.9	-30.9	-13.7	p .95		
224	31.91	20.5	18.7	16.8	15.2	15.5	13.2	- 8.0	- 1.4	p .99		
225	10.33	n.e.	5.6	2.2	3.9	6.1	12.3	+39.7	+ 1.7	n.s.		
GMU TOT/ AVG. 4/	882.96	23.9	26.1	23.6	23.3	22.4	21.9	- 3.1	- 0.8	p .99		

See footnotes at end of table, p. 36.



Table 3.--Trends in mourning dove breeding density indexes by physiographic region,  
1965-1970--continued

Region	Weight <sub>1</sub> / factor	ADJUSTED AVERAGE DOVES HEARD/ROUTE					2/3/		LINEAR REGRESSION, 1965-1970		
		1965	1966	1967	1968	1969	1970	Mean Annual Chg. (BYI)	Percent	No. of Doves	Stat. Sign. of Trend
WESTERN MANAGEMENT UNIT											
180	8.00	3.7	5.6	6.4	7.7	7.7	7.4	+12.7	+ 0.7		p .95
190	32.19	8.2	9.9	7.1	6.1	7.2	8.2	- 3.3	- 0.3		n.s.
201	27.15	8.9	9.9	9.0	11.2	9.1	8.3	- 1.0	- 0.1		n.s.
202	5.05	6.5	7.2	5.8	8.4	5.9	7.5	+ 1.6	+ 0.1		n.s.
203	19.43	6.8	16.0	21.1	16.4	17.3	18.6	+11.2	+ 1.7		n.s.
204	11.00	12.2	23.3	22.2	20.9	14.4	22.6	+ 3.6	+ 0.7		n.s.
205	6.17	92.8	172.3	39.8	30.9	44.3	28.7	-30.6	-20.4		n.s.
211	8.20	7.7	10.2	9.5	5.0	0.0	6.7	-17.1	- 1.1		n.s.
212	5.96	8.2	22.7	55.7	9.9	14.4	5.4	-12.1	- 2.4		n.s.
213	12.24	0.9	1.4	4.0	4.2	6.2	8.1	+89.5	+ 1.4		p .99
214	12.54	8.3	16.9	7.1	9.0	10.3	8.8	- 4.3	- 0.5		n.s.
215	13.78	6.9	11.8	13.7	10.0	29.2	14.4	+20.3	+ 2.5		n.s.
216	1.46	84.7	75.6	28.7	46.5	20.6	17.9	-30.9	-13.7		p .95
221	115.89	11.1	11.5	14.1	13.8	10.2	9.0	- 3.6	- 0.4		n.s.
222	35.86	50.0	49.5	36.8	31.1	27.6	18.7	-18.0	- 6.5		p .99
223	4.20	25.4	21.9	22.4	18.5	19.6	26.1	- 0.9	- 0.2		n.s.
224	24.20	20.5	18.7	16.8	15.2	15.5	13.2	- 8.0	- 1.4		p .99
231	7.99	16.9	12.1	11.9	12.2	15.2	11.2	- 4.0	- 0.5		n.s.
232	14.07	7.9	8.1	5.5	4.9	3.0	2.1	-24.2	- 1.3		p .99
233	4.42	46.6	20.0	19.0	No Estimate			n.e.	n.e.		n.e.
234	16.87	11.0	12.2	11.9	10.9	9.3	10.7	- 2.9	- 0.3		n.s.
241	8.84	0.5	0.0	2.5	2.5	0.8	0.8	+10.1	+ 0.1		n.s.
242	2.95	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		n.s.
243	8.29	28.2	10.1	4.0	28.2	2.0	6.0	-24.1	- 3.2		n.s.
244	10.48	6.6	5.4	4.9	6.8	7.3	7.1	+ 4.7	+ 0.3		n.s.
245	13.01	20.5	15.3	12.6	13.3	13.5	11.1	-10.0	- 1.5		p .95

See footnotes at end of table, p. 36.

Table 3.--Trends in mourning dove breeding density indexes by physiographic region, 1965-1970--continued

Region	Weight <sub>1</sub> / factor	ADJUSTED AVERAGE DOVES HEARD/ROUTE						LINEAR REGRESSION, 1965-1970		
		2/3/ 1970						Mean Annual Chg. (BYI)	Stat. Sign.	Percent No. of Doves of Trend
		1965	1966	1967	1968	1969	1970			
WESTERN MANAGEMENT UNIT (continued)										
246	19.64	31.2	26.2	22.8	23.5	21.3	21.1	- 7.4	- 1.8	p .95
247	8.10	7.4	5.2	4.3	1.8	6.0	6.0	- 3.9	- 0.2	n.s.
250	1.50	N o E s t i m a t e						n.e.	n.e.	n.e.
WMU TOT/ AVG. 4/	459.48	16.4	16.3	15.7	14.0	13.9	12.4	- 4.0	- 0.7	p .99

1/ Assigned physiographic region land-area value.

2/ From a Base Year, average doves heard calling per route for each physiographic region has been adjusted annually according to the percent change from the preceding year in counts on comparable routes. The base year represents the mean numbers of doves heard per route in 1968, 1969, and 1970; the resultant value being assigned to 1969.

3/ When a physiographic region is bisected by a management unit boundary, its mean breeding density index is assigned to both units.

4/ The management unit breeding density indexes are obtained from Table 2.



As the Nation's principal conservation agency, the Department of the Interior has basic responsibilities for water, fish, wildlife, mineral, land, park, and recreational resources. Indian and Territorial affairs are other major concerns of this department of natural resources.

The Department works to assure the wisest choice in managing all our resources so that each shall make its full contribution to a better United States now and in the future.



---

UNITED STATES  
DEPARTMENT OF THE INTERIOR  
FISH AND WILDLIFE SERVICE  
BUREAU OF SPORT FISHERIES AND WILDLIFE  
WASHINGTON, D. C. 20240

POSTAGE AND FEES PAID  
U.S. DEPARTMENT OF THE INTERIOR

